arXiv:hep-th/0410159v3 22 Dec 2004

Extended Non Linear Conformal Symmetry and DSR Velocities on the Physical Surface

Carlos Leiva*
Departamento de Física, Universidad de Tarapacá
Casilla 7-D, Arica, Chile

(Dated: February 1, 2008)

The relation between Conformal generators and Magueijo Smolin Deformed Special Relativity term, added to Lorentz boosts, is achieved. The same is performed for Fock Lorentz transformations. Through a dimensional reduction procedure, it is demonstrated that a massless relativistic particle living in a d dimensional space, is isomorphic to one living in a $d+2$ space with pure Lorentz invariance and to a particle living in a AdS_{d+1} space. To accomplish these identifications, the Conformal Group is extended and a nonlinear algebra arises. Finally, because the relation between momenta and velocities is known, the problem of position space dynamics is solved.

PACS numbers: PACS Nos.: 02.20.Hj / 03.30.+p

I. INTRODUCTION

It has been claimed that the special relativity must be modified, that a Lorentz symmetry breaking could be observable in the future in high energy cosmic ray spectra [1, 2], and corrections in the dispersion relation $E^2-p^2=m^2$ has been proposed [3]. Furthermore, quantum gravity models suggest that it could be desirable to review the Lorentz invariance relations and string theories consider some modifications to the very structure of the space time at high energy scales. Some data seem to invite to introduce a minimal length in physical theories, indeed, there are theories that have some fundamental quantities: the Planck longitude $l_p=\sqrt{\hbar G/c^3}$, its associated time scale $t_p=l_p/c$ and the Planck energy $E_p=\hbar/t_p$. It is supposed that beyond these thresholds, the physics should change dramatically. However, these absolute values of longitude, time or energy are not in agreement with the Lorentz transformations, this point reinforces the idea of to modify the Lorentz boosts.

Solutions for the problem of how must be, the Lorentz boosts, modified has been proposed, a very interesting one is Doubly (or Deformed), special relativity (DSR) theories [4–6]. These theories are based on a generalization of Lorentz transformations through the more broad point of view of conformal transformatios, hey have two observer independent scales, velocity of light and Planck length.

Usually a modification of Lorentz boosts in momentum space is performed, however when this is done, retrieving the position space dynamics can be a highly problematic task due to the loss of linearity. Kimberly, Magueijo and Medeiros [7], have proposed some methods to achieve this, using a free field theory, other approaches can be seen in [8], specially the approach of Deriglazov is very interesting because he starts from a conformal group, but it is different from the one proposed here, because he start at the position space [9]. So, this is a worth investigating aspect of DSR theories that is not so understood until today. Other approaches have been done in order to identify AdS spaces as arenas for DSR theories and the approach of this paper can add some information in that way too.

DSR theories are of increasing interest because they can be useful as a new tool in gravity theories, in Cosmology as an alternative to inflation [10, 11], and in other fields like propagation of light [12], that is related, for instance, to cosmic microwave background radiation.

In this paper it is demonstrated that it is possible, formally, to obtain Fock-Lorentz [13, 14]and Magueijo-Smolin deformations, trough a reduction process and using the conformal group generators as the generators of the deformed Lorentz algebra. Then, the open problem of obtaining the space time dynamics is solved through the relationship on the physical surface between momenta and velocities on the physical surface that rise up from the method of dimensional reduction.

More specifically, it is conjectured that the deformations of the Lorentz algebra in the Fock-Lorentz formulation, can be treated as a transformation made by a linear combination of conformal group generators and the momentum case (the Magueijo Smolin case), can be understood as the same process, but the inclusion of a new generator is needed. This new generator can be constructed from the same theory, and complete the set of symmetries of the massless Klein Gordon Equation. Then, a DSR massless particle is shown to be isomorphic to a normal Lorentz particle living in a $d+2$ space, and the deformations are induced by the dimensional reduction.

---

*cleivas@uta.cl

II. CONFORMAL SYMMETRY OF MASSLESS PARTICLE

The infinitesimal transformations

$$\delta x^\mu = \omega^\mu{}_\nu x^\nu + \alpha^\mu + \beta x^\mu + 2(x\gamma)x^\mu - x^2\gamma^\mu \tag{1}$$

generate the conformal symmetry, $ds^2 \to ds'^2 = e^{2\sigma}ds^2$, on the d-dimensional Minkowski space $\mathbb{R}^{1,d-1}$ with metric

$$ds^2 = dx^\mu dx^\nu \eta_{\mu\nu} = -dx_0^2 + \sum_{i=1}^{d-1} dx_i^2.$$

Here the parameters $\omega^\mu{}_\nu$, α^μ, β and γ^μ correspond to the Lorentz rotations, space-time translations, scale and special conformal transformations. Due to a nonlinear (quadratic) in x^μ nature of the two last terms in (1), the finite version of the special conformal transformations,

$$x'^\mu = \frac{x^\mu - \alpha^\mu x^2}{1 - 2\alpha x + \alpha^2 x^2}, \tag{2}$$

is not defined globally, and to be well defined requires a compactification of $\mathbb{R}^{1,d-1}$ by including the points at infinity.

On the classical phase space with canonical Poisson bracket relations $\{x_\mu, p_\nu\} = \eta_{\mu\nu}$, $\{x_\mu, x_\nu\} = \{p_\mu, p_\nu\} = 0$, the transformations (1) are generated by

$$\begin{aligned} M_{\mu\nu} &= x_\mu p_\nu - x_\nu p_\mu, \\ P_\mu = p_\mu, &\qquad D = x^\mu p_\mu, \\ K_\mu &= 2x_\mu(xp) - x^2 p_\mu. \end{aligned} \tag{3}$$

The generators (3) form the conformal algebra

$$\begin{aligned} \{M_{\mu\nu}, M_{\sigma\lambda}\} &= \eta_{\mu\sigma}M_{\nu\lambda} - \eta_{\nu\sigma}M_{\mu\lambda} \\ &\quad + \eta_{\mu\lambda}M_{\sigma\nu} - \eta_{\nu\lambda}M_{\sigma\mu}, \\ \{M_{\mu\nu}, P_\lambda\} &= \eta_{\mu\lambda}P_\nu - \eta_{\nu\lambda}P_\mu, \\ \{M_{\mu\nu}, K_\lambda\} &= \eta_{\mu\lambda}K_\nu - \eta_{\nu\lambda}K_\mu, \\ \{D, P_\mu\} &= P_\mu, \\ \{D, K_\mu\} &= -K_\mu, \\ \{K_\mu, P_\nu\} &= 2(\eta_{\mu\nu}D + M_{\mu\nu}), \\ \{D, M_{\mu\nu}\} &= \{P_\mu, P_\nu\} = 0, \\ \{K_\mu, K_\nu\} &= 0 \end{aligned} \tag{4}$$

The algebra (4) is isomorphic to the algebra $so(2,d)$, and by defining

$$\begin{aligned} J_{\mu\nu} = M_{\mu\nu}, &\quad J_{\mu d} = \frac{1}{2}(P_\mu + K_\mu), \\ J_{\mu(d+1)} = \frac{1}{2}(P_\mu - K_\mu), &\quad J_{d(d+1)} = D, \end{aligned} \tag{5}$$

can be put in the standard form

$$\begin{aligned} \{J_{AB}, J_{LN}\} &= \eta_{AL}J_{BN} - \eta_{BL}J_{AN} \\ &\quad + \eta_{AN}J_{LB} - \eta_{BN}J_{LA} \end{aligned} \tag{6}$$

with $A, B = 0, 1, d, d+1$, and

$$\eta_{AB} = diag(-1, +1, \ldots, +1, -1). \tag{7}$$

The phase space constraint $\varphi_m \equiv p^2 + m^2 = 0$ describing the free relativistic particle of mass m in $\mathbb{R}^{1,d-1}$ is invariant under the Poincaré transformations, $\{M_{\mu\nu}, \varphi_m\} = \{P_\mu, \varphi_m\} = 0$. Unlike the P_μ and $M_{\mu\nu}$, the generators of the scale and special conformal transformations commute weakly with φ_m only in the massless case $m = 0$: $\{D, \varphi_0\} = 2\varphi_0 = 0$, $\{K_\mu, \varphi_0\} = 4x_\mu\varphi_0 = 0$.

III. DIMENSIONAL REDUCTION

It was demonstrated in Leiva and Plyushchay [15], that a dimensional reduction process from a massless particle living in $(d+2)$ dimensional space $\mathbb{R}^{2,d}$, with coordinates H^A and metric (7), and introducing canonical momenta Π_A, ($\{H_A, \Pi_B\} = \eta_{AB}$) can transform the $so(2,d)$ generators

$$\mathfrak{J}_{AB} = H_A\Pi_B - H_B\Pi_A., \tag{8}$$

into the conformal generators in (d)-dimensional space $\mathbb{R}^{1,d-1}$. In order to achieve this, three constrains are added to the theory:

$$\phi_0 \equiv \Pi_A\Pi^A = 0, \tag{9}$$

$$\phi_1 \equiv H^A H_A = 0, \qquad \phi_2 \equiv H^A\Pi_A = 0. \tag{10}$$

The process is implemented trough suitable canonical transformations and then a dimensional reduction is performed onto the surface defined by ϕ_1 and ϕ_2. So in terms of new $\mathbb{R}^{2,d}$ canonical variables $\tilde H^\mu$, $\tilde\Pi_\mu$ with $\{\tilde H^\mu, \tilde\Pi_\mu\} = 1$ as defined in [15], the $so(2,d)$ generators are the following:

$$\begin{aligned} \mathfrak{J}_{\mu\nu} &= \tilde H_\mu\tilde\Pi_\nu - \tilde H_\nu\tilde\Pi_\mu, \\ \mathfrak{J}_{\mu+} &= \tilde\Pi_\mu, \\ \mathfrak{J}_{d(d+1)} &= \tilde H_\mu\tilde\Pi^\mu + 2\frac{\tilde\Pi_+}{\tilde H^-}\phi_1 - \phi_2, \\ \mathfrak{J}_{\mu-} &= 2(\tilde H_\nu\tilde\Pi^\nu)\tilde H_\mu - (\tilde\Pi_\nu\tilde H^\nu)\tilde\Pi_\mu \\ &\quad + \frac{1}{\tilde H^{-2}}(\tilde\Pi_\mu + 4\tilde\Pi_+\tilde H^-\tilde H_\mu)\phi_1 - 2\tilde H_\mu\phi_2. \end{aligned}$$

To execute this process it is supposed that $\tilde H^- \neq 0$. The constraints now, look like:

$$\tilde\Pi_\mu\tilde\Pi^\mu = 0, \tag{11}$$

$$\tilde H^+ = 0, \qquad \tilde\Pi_- = 0. \tag{12}$$

The variables $\tilde\Pi_\mu$ and $\tilde H_\mu$ are observable (gauge invariant) variables with respect to the constraints (11) and

(12), whereas $\tilde{H}^{-} = H^{-}$ and $\tilde{\Pi}_{+} = \Pi_{+}$ are not. They can be removed by introducing the constraints

$$\phi_3 \equiv H^{-} + 1 = 0, \quad \phi_4 \equiv \Pi_{+} = 0 \tag{13}$$

as gauge fixings of (12) and the reduction to the physical surface achieves the conformal generators in $\mathbb{R}^{1,d-1}$ as projections of the $so(2,d)$ generators. Now, with the identification $x^\mu = \tilde{H}^\mu$ and $p_\mu = \tilde{\Pi}_\mu$ and (11), the original massless relativistic particle is totally retrieved.

IV. EXTENDED NON-LINEAR CONFORMAL ALGEBRA

It is possible, trough the procedure described in the last section, and because positions and momenta are at same level in the hamiltonian formulation, to introduce a new generator $\tilde{K}_\mu$:

$$\tilde{K}_\mu = 2p_\mu(xp) - p^2 x_\mu \tag{14}$$

To do this, we can perform a canonical transformation in (11), $\tilde{H}^A \to -\breve{\Pi}^A$ and $\tilde{\Pi}^A \to \breve{H}^A$ and now to identify $x^\mu = \breve{H}_\mu$ and $p_\mu = \breve{\Pi}_\mu$. Doing this, $\mathfrak{J}_{\mu-}$ is projected as $\tilde{K}$, and $\mathfrak{J}_{\mu+}$ is projected as x^μ, but the particle has now the constrain $x^2 = 0$. The nature of this generator is totally new because it produces conformal transformations on the momentum space and the finite version of this transformations is :

$$p'^\mu = -\frac{p^\mu - \alpha^\mu p^2}{1 - 2\alpha p + \alpha^2 p^2}, \tag{15}$$

This is is a symmetry of the massless Klein-Gordon equation too, in fact: $\{\tilde{K}, P^2\} = 2P^2 P_\mu = 0$. So, after the inclusion of $\tilde{K}$, aside the relations (4) we have:

$$\begin{aligned}
\{M_{\mu\nu}, \tilde{K}_\lambda\} &= \eta_{\mu\lambda}\tilde{K}_\nu - \eta_{\nu\lambda}\tilde{K}_\mu, \\
\{D, \tilde{K}_\mu\} &= \tilde{K}_\mu, \\
\{\tilde{K}_\mu, P_\nu\} &= 2P_\mu P_\nu - P^2\eta_{\mu\nu}, \\
\{\tilde{K}_\mu, \tilde{K}_\nu\} &= 0, \\
\{\tilde{K}_\mu, K_\nu\} &= 4(M_{\mu\nu} - D\eta_{\mu\nu})D - 2P_\mu K_\nu \\
&\quad + \frac{2(2DP_\mu - \tilde{K}_\mu)(2DP_\nu - \tilde{K}_\nu)}{\tilde{K}P}
\end{aligned} \tag{16}$$

Due to the last bracket, the set of generators form a non linear algebra.

We can then, to have two separate set of symmetries. Each set form a dynamic $SO(2,1)$ symmetry of the Klein Gordon equation,: one that includes K_μ, generating special conformal transformations in the position space. And the other that includes $\tilde{K}_\mu$, generating special conformal transformations in the momentum space.

The existence of $\tilde{K}_\mu$ could produce some consequences in the symmetry group of non relativistic particles since there is a canonical relation between a non relativistic particle in $\mathbb{R}^1$ and the relativistic one in $\mathbb{R}^{1,1}$ [15]. This can be of some interest because Fluid Mechanics symmetries are intimately related to the free non relativistic particle symmetry [16].

Each group of $SO(2,1)$ symmetries can be used to construct the DSR transformations in position or momentum space. In the two next sections, both cases shall be reviewed

V. FOCK-LORENTZ TRANSFORMATIONS

The Fock-Lorentz transformation [13] [14], are introduced as general linear-fractional transformations between spatial and temporal coordinates and the result is leaded by symmetry arguments to the final form:

$$\begin{aligned}
t' &= \frac{\gamma(u)(t - \mathbf{ur}/c^2)}{1 - (\gamma(u) - 1)ct/R + \gamma(u)\mathbf{ur}/Rc}, && (17) \\
\mathbf{r}\| &= \frac{\gamma(u)(\mathbf{r}\| - \mathbf{u}t)}{1 - (\gamma(u) - 1)ct/R + \gamma(u)\mathbf{ur}/Rc}, && (18) \\
\mathbf{r}\perp &= \frac{\mathbf{r}\perp}{1 - (\gamma(u) - 1)ct/R + \gamma(u)\mathbf{ur}/Rc}, && (19) \\
& && (20)
\end{aligned}$$

where $\mathbf{r}\|$ is the component of the position vector parallel to the boost $\mathbf{u}$ and $\mathbf{r}\perp$ is the perpendicular one. Where γ is the Lorentz factor $\sqrt{1 - u^2/c^2}$ and R is a constant with the dimension of length.

These transformations can be seen as Lorentz transformations for the quantities:

$$\tilde{t} = \frac{t}{1 + ct/R}, \qquad \tilde{\mathbf{r}} = \frac{\mathbf{r}}{1 + ct/R} \tag{21}$$

They can be treated as results of a infinitesimal transformations:

$$\delta x^\mu = \alpha^\nu\{x^\mu, 2x_\nu(xp)\} \tag{22}$$

where $x^0 = ct$.

In order to obtain this generator, it can be represented as the projection of

$$(\mathfrak{J}_{\mu-} + \mathfrak{J}_{\mu+}\mathfrak{J}_{d(d+1)}), \tag{23}$$

in R^{d+2} space, wich is a nonlinear combination of the isometries there. However $2x_\nu(xp)$ is not a symmetry of the Klein-Gordon equation because $\{2x_\nu(xp), \varphi_0\} = 4p_\mu(xp) + 4x_\mu\varphi_0 \neq 0$. So, it is not possible to include it into the generators set of the massless relativistic particle symmetries.

An alternative, and linear solution is to see (21) as (2) reduced to the surface $x^2 = 0$ and with the choice $\alpha = (-1/2R, 0, 0, 0)$ and identifying x^0 with ct. In this sense, the Fock Lorentz transformations can be seen as a linear combination of conformal generators

$$M_{0\mu} + K_\mu,$$

that is a projection of

$$(\mathfrak{J}_{\mu\nu} + \mathfrak{J}_{\mu-})$$

(which is linear too), for a particle living on the cone (light cone) $x_1^2 + x_2^2 + x_3^2 = x_0^2$.

Then we can construct the Fock-Lorentz transformations for a particle living on the light cone through the projection of the action of a linear combination of conformal generators.

Then, Fock Lorentz appear to be deformations of the usual Lorentz isometries induced by a projection of the system on the physical surface.

VI. THE MAGUEIJO SMOLIN DSR MOMENTUM TRANSFORMATION

On the other hand, the Lorentz boost proposed in Magueijo and Smolin [4]

$$K^i = L_{0i} + l_p p^i p_\mu \frac{\partial}{\partial p_\mu} \tag{24}$$

where l_p is the Planck length, can be exponentiated as

$$K^i = U^{-1}(p_0) L_{0i} U(p_0) \tag{25}$$

where $U(p_0) = exp(l_p p_0 p_\mu \frac{\partial}{\partial p_\mu})$, and the action of $U(p_0)$ over p_μ is

$$U(p_0) p_\mu = \frac{p_\mu}{1 - l_p p_0} \tag{26}$$

The generator added to the Lorentz boost in (29) is the operational representation of $p^i(xp)$ and it can be as $\tilde{K}_\mu$, that produces the following transformation :

$$p'^\mu = -\frac{p^\mu - \alpha^\mu p^2}{1 - 2\alpha p + \alpha^2 p^2}, \tag{27}$$

to achieve the identification with (31), the transformation parameter must be $\alpha^\mu = (-l_p/2, 0, 0, 0)$ and the global minus sign must be removed.

It is possible then, to prescribe how to obtain the position and momenta transformation using conformal generators with some $\tilde{K}$ added. Where does this new generator come from? Some clues will be seen in the next section.

Starting with the ordinary Lorentz generators:

$$L_{ab} = p_a \frac{\partial}{\partial p^b} - p_b \frac{\partial}{\partial p^a} \tag{28}$$

The modified Lorentz boost proposed in Magueijo and Smolin [4] is:

$$K_i = L_{0i} + l_p p_i p_\mu \frac{\partial}{\partial p_\mu}, \tag{29}$$

where the second term is the deformation proposed and l_p is the Planck length. It can be exponentiated as:

$$K^i = U^{-1}(p_0) L_{0i} U(p_0) \tag{30}$$

where $U(p_0) = exp(l_p p_0 p_\mu \frac{\partial}{\partial p_\mu})$, and the action of $U(p_0)$ over p_μ is

$$U(p_0) p_\mu = \frac{p_\mu}{1 - l_p p_0} \tag{31}$$

Following the Magueijo Smolin procedure, it can be seen that boosts in the z direction (as an example), are :

$$\begin{aligned} p_0' &= \frac{\gamma(p_0 - v p_z)}{1 + l_p(\gamma - 1)p_0 - l_p \gamma v p_z} \\ p_z' &= \frac{\gamma(p_z - v p_0)}{1 + l_p(\gamma - 1)p_0 - l_p \gamma v p_z} \\ p_x' &= \frac{p_x)}{1 + l_p(\gamma - 1)p_0 - l_p \gamma v p_z} \\ p_y' &= \frac{p_y)}{1 + l_p(\gamma - 1)p_0 - l_p \gamma v p_z}. \end{aligned} \tag{32}$$

This transformations are identical to those obtained by Fock [13, 14], but applied to momentum space. They can be retrieved replacing p by x and p_0 by t, instead.

The Fock like transformations can also be obtained as usual Lorentz transformations for the transformed p' from eq 31 [13, 14], so we need lead just with the extra term of the deformed boosts.

The transformation (31) can be yielded as result of the action of an $\tilde{K}_\mu = 2p_\mu(xp) - p^2 x_\mu$ generator, that produces:

$$p'^\mu = -\frac{p^\mu - \alpha^\mu p^2}{1 - 2\alpha p + \alpha^2 p^2}. \tag{33}$$

So, the Magueijo Smolin transformation can be seen as a combination of

$$-M_{0\mu} + \tilde{K}_\mu,$$

that is a projection of

$$(\mathfrak{J}_{\mu\nu} + \mathfrak{J}_{\mu-})$$

as projected from the R^{d+2} to the particle living on the light cone, $x_1^2 + x_2^2 + x_3^2 = x_0^2$, this is the same condition we had before, and it is a condition for a truly massless relativistic particle.

Then we can construct the Magueijo Smolin transformations for a particle living on the light cone through the projection of the action of a linear combination of conformal generators. But we must consider a massless particle, then we can achieve the identification with (31), in order to do that it is necessary to transform the constrain $\psi_1 = p^2 \approx 0$ in an exact identity, the transformation parameter must be $\alpha^\mu = (l_p/2, 0, 0, 0)$ and the global minus sign must be removed.

VII. THE PROJECTION ONTO THE PHYSICAL SURFACE.

The fundamental problem with the momentum space formulation is to retrieve the position space. This is a task that has called the attention of many researches [7–9]. The advantage of the point of view in the way that is managed here, is that the task is replaced by the known problem of relating momenta and velocities when a dimensional reduction is performed. Just as an example, let's recall the process to project a free massless relativistic particle onto the physical surface. This particle has the Lagrangian:

$$L = \frac{\dot{x}^2}{2e} \tag{34}$$

the constrain that rules the dynamic of the particle are:

$$\varphi_e = p_e \approx 0 \tag{35}$$

$$\varphi_1 = \frac{1}{2}(p^2) \approx 0 \tag{36}$$

The condition over the conjugated momentum of e, $p_e = 0$ shows that e is a non dynamic variable and we will not take care of it in the following.

Due the constrain φ_1, the relation between momenta in phase space and velocities in the configuration space is not so clear as it would be desirable, because all momenta yielding Eq. (36) should be identified. This ambiguity can be seen as produced by the fact that the system is constrained to a subspace in the phase space, this subspace is usually named as the physical surface.

In order to obtain a truly relationship between momenta and velocities a reduction of the system to this physical surface must be performed, this can be done by the following process:

First, we can introduce a gauge fixing to the constrain Eq. (36), a suitable one is:

$$\varphi_2 = x^0 - \tau \approx 0 \tag{37}$$

Now, it is necessary to reformulate the parenthesis definition abandoning the Poisson definition and using the Dirac Parenthesis instead. It is defined by the relation:

$$\{A, B\}_D = \{A, B\}_P - \{A, \varphi_i\}_P C_{ij}^{-1} \{\varphi_j, B\}_P, \tag{38}$$

where $\{\}_P$ indicates Poisson bracket and $\{\}_D$ Dirac bracket. The matrix C is defined by:

$$C_{ij} = \{\varphi_i, \varphi_j\}_P \tag{39}$$

in this case:

$$C_{i,j} = p_0 \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}, \tag{40}$$

with

$$C_{i,j}^{-1} = \frac{1}{p_0} \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \tag{41}$$

So, the Dirac brackets are:

$$\{x^0, p_0\}_D = 0 \tag{42}$$

$$\{x^0, p_i\}_D = \frac{p_i}{p^0} \tag{43}$$

$$\{x^i, p^j\}_D = \delta^{ij} \tag{44}$$

Now, in order to retrieve the Hamiltonian relations, we define the independent variables:

$$u^a = (x^i, p_i) \tag{45}$$

and introduce the following parameterization:

$$z^s(u, \tau) = (x^0 = \tau, x^i, p^0 = +\sqrt{p_i^2}, p^i) \tag{46}$$

It is, now, necessary to redefine our Hamiltonian, this can be performed setting:

$$\mathcal{H} = H + F$$

where H is the former canonical Hamiltonian and F is a correction due to the projection of the system to the physical surface, F yields the following equation:

$$\frac{\partial F}{\partial u^m} = -\frac{\partial z^s}{\partial u^m} W_{st} \frac{\partial z^t}{\partial \tau} \tag{47}$$

where W_{st} is defined by:

$$W_{st} = \{x^\mu, p_\nu\}$$

$$W_{st} = \begin{pmatrix} 0 & -\delta^{\mu}_{\nu} \\ \delta^{\mu}_{\nu} & 0 \end{pmatrix}, \tag{48}$$

after proper calculations we have

$$F = \sqrt{\vec{p}^2} \tag{49}$$
$$H = 0 \tag{50}$$
$$\mathcal{H} = \sqrt{\vec{p}^2} \tag{51}$$

and the velocity induced on the physical surface is:

$$\dot{x}^i(\tau) = \{x^i, \mathcal{H}\}_D = \frac{p^i}{\sqrt{\vec{p}^2}} \tag{52}$$

Just now, we are sure what the relation between velocities and momenta is, on the physical surface and we can perform the transformation (33):

$$\bar{p}_i = \frac{p_i}{1 - 2l_p\sqrt{p_i^2}} \tag{53}$$

and, identifying $\sqrt{p_i^2}$ with the energy E, the velocity expression yields:

$$\begin{aligned} \dot{x}^i &= \frac{p^i}{E} \\ \dot{\bar{x}}^i &= \frac{p^i}{E - 2l_pE^2} \end{aligned} \tag{54}$$

Now, in order to obtain the complete Magueijo Smolin velocity we just need to perform an usual Lorentz boost, for example in z direction:

$$\begin{aligned} v_x &= \frac{(1-u^2)P_x}{(E - 2l_pE^2 - uP_z)} \\ v_y &= \frac{(1-u^2)P_y}{(E - 2l_pE^2 - uP_z)} \\ v_x &= \frac{P_z - u(E - 2l_pE^2)}{(E - 2l_pE^2 - uP_z)} \end{aligned} \tag{55}$$

VIII. DSR AND ADS SPACES

The $(d+2)$-dimensional system (9), (10) can be reinterpreted as a massless particle living on the border of the AdS$_{d+1}$ of infinite radius, whose isometry corresponds to the conformal symmetry of the massless particle in d dimensions. This can be done in the following way. The massless particle on AdS$_{d+1}$ of radius R can be described by the constraints

$$\phi_0 = \mathfrak{P}_A\mathfrak{P}^A = 0, \qquad \phi_1 = \mathfrak{X}_A\mathfrak{X}^A + R^2 = 0, \tag{56}$$

where $\mathfrak{X}^A$ and $\mathfrak{P}_A$ are the canonical variables. The Poisson brackets of the constraints (56) are $\{\phi_1, \phi_0\} = 4\mathfrak{X}^A\mathfrak{P}_A$. To have the reparametrization-invariant system, the ϕ_0 has to be the first class constraint. This can be achieved by postulating the constraint

$$\phi_2 = \mathfrak{X}^A\mathfrak{P}_A = 0 \tag{57}$$

in addition to the constraints (56). The constraint (57) generates the local scale transformations which due to the relation $\{\phi_2, \phi_0\} = 2\phi_0$ are consistent with the reparametrization invariance generated by the constraint ϕ_0. Since $\{\phi_2, \phi_1\} = -\phi_1 + R^2 \neq 0$, the constraint $\phi_1 = 0$ can be understood as a gauge condition for the constraint (57). Now, let us realize a canonical transformation $(\mathfrak{X}^A, \mathfrak{P}_A) \rightarrow (X^A, P_A)$,

$$X^A = \frac{\mathfrak{X}^A}{R^{1+\varepsilon}}, \quad P_A = \mathfrak{P}_A R^{1+\varepsilon}, \tag{58}$$

with a constant $\varepsilon > 0$, and take a limit $R \rightarrow \infty$, $\mathfrak{X}^A \rightarrow \infty$, $\mathfrak{P}_A \rightarrow 0$ in such a way that the variables (58) would be finite. Then, the constraints (56), (57) take the form of the first class constraints (9), (10), and we reproduce the system on the cone. Because of the change of the nature of the constraints from the second to the first class, the described limit procedure has a rather formal character, however a relationship between DSR transformations and the symmetries of a AdS$_{d+1}$ of infinite radius can be observed, because the DSR particle in d is isomorphic to one moving in AdS$_{d+1}$ with Lorentz invariance.

IX. DISCUSSION AND OUTLOOK

To conclude, let us summarize the obtained results and discuss shortly some problems that deserve a further attention.

In section 2, the Conformal Group of the massless relativistic particle was reviewed and in section 3, it was obtained as a dimensional reduction from a $\mathbb{R}^{d+2}$ space. In section 4, Because of the interchangeability of momenta and positions in the Hamiltonian formulation, a new generator was added. With this generator, the Conformal Group can be extended and a non linear algebra rose. In section 4, starting with the formulation of Magueijo Smolin deformed boosts, they are realized as a version of the special conformal generator after a dimensional reduction onto the mass shell of a massless relativistic particle and then, with the Hamiltonian relations retrieved on that physical surface, the velocity can be achieved from the transformed momenta.

Among the problems that deserve more attention is the problem of how to include the mass term, that is still in darkness. On the other hand, due the relation of DSR theories with AdS spaces concluded here and in the bibliography, it could be interesting to research about its relation with the AdS/CFT correspondence. Finally to extend the studies to fields formulation could be a interesting task too.

Acknowledgements

I want to thank to the Departamento de Física de la Universidad de Tarapacá.

[1] G. Amelino-Camelia and Tsvi Piran, *Planck-scale deformation of Lorentz symmetry as a solution to the UHECR and $Tev - \gamma$ paradoxes*, Phys. Rev.D64(2001) 036005 [arXiv:astro-ph/0008107].

[2] F.W. Stecker and S.T. Scully, *Lorentz Invariance Violation and the Spectrum and Source Power of Ultrahigh Energy Cosmic Rays*, [arXiv:astro-ph/0412495]

[3] G. Amelino-Camelia and T. Piran, Phys. Rev. D64 (2001) 036005.

[4] J. Magueijo and L. Smolin, Phys. Rev. Lett. **88**, 190403(2002) [arXiv:hep-th/0112090]

[5] G. Amelino-Camelia, Int. J. Mod. Phys. D11, 35, 2002, [arXiv:gr-qc/0012051], Ohys. Lett. B510, 255-263, 2001.

[6] J. Kowalski-Glikman, Phys. Lett. A286, 391-394, 2001, [arXiv:hep-th/0102098]; N.R. Bruno, G. Amelino-Camelia, J. Kowalski-Glikman, Phys. Lett. B522, 133-138,2001.

[7] D. Kimberly, J. Magueijo and J. Medeiros [arXiv:gr-qc/0303067]

[8] Sijie Gao and Xiaoning Wu, *Position space of Doubly Special Relativity* [arXiv:gr-qc/0311009]

[9] A. A. Deriglazov, Doubly special relativity in position space starting from the conformal group, hep-th/0409232.

[10] J. Moffat, Int. J. of Physics D **2**, 351 (1993); J. Moffat, Foundations of Physics, **23**, 411 (1993)

[11] A. Albrecht and J. Magueijo, Phys. Rev. D **59**, 043516 (1999)

[12] Sung Ku Kim, Sun Myong Kim, Chaiho Rim and Jae Hyung Yee, *Propagation of Light in Doubly Special Relativity* [arXiv:gr-qc/0401078]

[13] V. Fock, *The theory of space-time and gravitation*, Pergamon Press,1964.

[14] S.N. Manida, [arXiv:gr-qc/9905046]

[15] C. Leiva and M. S. Plyushchay, *Conformal Symmetry of Relativistic and Nonrelativistic Systems and AdS/CFT Correspondence*, Annals of Physics 307 0310 (2003) 372-391. [arXiv:hep-th/0301244].

[16] L. O'raifeartaigh and V. V. Sreedhar, Annals Phys. 293 (2001) 215 [arXivhep-th/0007199]